



Physicochemical characterization, fatty acid in Corn Seed Oil using GC-FID method

Purendra Singh

Department of Chemistry, Banasthali Vidyapith, Rajasthan, India

Abstract

Physical properties like physical state at 25^o C/ 1Atm, colour, odour, solubility, miscibility and physicochemical properties like density /specific gravity, viscosity, flash point, boiling point, saponification value, unsaponifiable value, iodine value, colour (Lovibond tintometer), peroxide value and refractive Index Corn oil were studied to evaluate the compositional quality of oils and also to investigate the effect on the use of same oil for repeated frying as it ultimately changes the nutritional, physicochemical, and sensory properties of the oil. The physicochemical analysis showed color values were 3.0 red maximum, density (g/mL)/specific gravity were 0.918 at 25^oC, viscosity (millipoise) were 114.00, flash point were 305, boiling point were 148 ^oC, saponification value (mg) were 153.6, unsaponifiable value (gm) were 2.2, iodine value (g) were 15.77, peroxide value (meq/kg) were 0.165, refractive index (40^o C) were 1.466. The analyses of fatty acid profile showed higher linoleic acid, oleic acid and palmitic acid contents and saturated fatty acid were 15.282, unsaturated fatty acid were 84.717 present in corn oil.

Keywords: Corn seed, Corn seed oil, Fatty Acid Profile, Quality characteristics

Introduction

Nuts are a good source of oil containing higher unsaturated fatty acids (PUFA and MUFA) to saturated fatty acids ratio the higher body weight is associated with higher percentage of fat in the diet [18, 19]. Fats and oils consist of a wide group of compounds that are soluble in organic solvents and insoluble in water. They have lower densities than water and at normal room temperature range inconsistently from liquids to solids depending on their structure and composition. The words oils, fats, and lipids are all used to refer to fat; oils are usually used to refer to fats that are liquids at room temperature, while fats are usually used to refer to that are solid at normal temperature [20, 21].

In quality control of oils, several parameters such as saponification value (average molecular weight), iodine value (degree of unsaturation), unsaponifiable value and peroxide value as well as the fatty acid profile are of interest as they determine the quality and hence the economic value of the product. Currently, the majority of these parameters are determined by using classical wet chemical methods [5]. Edible oils have been analyzed for different metals using atomic absorption spectrometer (FAAS and GFAAS), inductively coupled plasma optical emission spectrometer (ICP-OES), and inductively coupled plasma mass spectrometer (ICP-MS). Atomic absorption spectroscopy (flame, graphite furnace, hydride generation, and cold vapour) is still the most widely used technique [21-24].

Maize (*Zea mays* L.) has been a crop extended and an important food for the American societies since pre-hispanic times [1].

Maize (*Z. mays* L.) is the third most important food crop in the world and a major source of energy, protein, and other nutrients for both human and livestock. Maize contains 7-13 g/100 g proteins (d.m.). However, the quality of maize proteins is poor, because they are deficient in the essential amino acids lysine and tryptophan [2, 3].

Corn is successful cereal grass useful as food, feed and in industries. Corn is also source of corn oil which is tasteless odourless and pale yellow oil derived from the corn kernel. Normal corn has usually oil ranging about 3-4 per cent oil, especially in the embryo. Corn with oil content more than six per cent is called high oil corn (HOC) [4, 5, 6, 7, 8, and 9].

Materials and Methods

Material and chemicals

Methanol, Hydrochloric acid, sodium hydroxide, potassium hydroxide, benzene, sodium thiosulphate, petroleum ether, potassium iodide (KI), hexane, acetic acid and chloroform were of A.R Grade and purchased from Merck Darmstadt Germany. Corn oils were extracted from their dried, dehulled and powdered seeds purchased from the local market in Alwar, Rajasthan (India).

Oil extraction

Corn is cultivated in the Alwar (Rajasthan). Corn oil sample was obtained from corn seeds using the cold pressed method. Oil was then stored at 4.0±2^oC. Oil extraction was conducted using a Soxhlet apparatus for approximately 6 hrs with hexane as solvent, with a solid to solvent ratio of 1/7 m/v. After the extraction process, the flask contents were filtered, and the liquid fraction containing the lipid extract and solvent was poured into a 500 mL flask of a rotary film evaporator to remove the solvent. The obtained oil was collected, evaporated under nitrogen, weighed, and stored in sealed amber glass vials at -20^oC until analysis [10]. Determination of physical properties such as physical state at 25^o C/ 1Atm, colour, odour, solubility and miscibility was determined by the manual of methods of analysis of food, fssai, 2015 [13].

Determination of physicochemical properties such as colour by using lovibond tintometer, flash point, unsaponifiable value was

determined by hand book of phytochemical, pharmacognostical, water and edible oil analysis (A lab manual) [17].

Density measurement

Densities of oil samples before and after frying were measured by an R.D bottle with a capacity of 10 mL [17].

Boiling point measurement

The boiling point of oil samples were measured by a thermometer. The boiling point depends upon the degree of unsaturation of fatty acids.

Viscosity measurement

The viscosity of oil samples before and after frying was measured by Viscometer techniconominal constant 0.05 Cs/c. The flow time of corn oil sample were recorded with a stop watch least count ± 0.01 s [17].

Frying process

Potatoes were peeled and cut into pieces (approx 18) and were fried in the frying oil at constant temperature at four frying times. Frying experiments were conducted in duplicate in each frying medium.

Saponification value

At first, 2 g oil was taken in 250 ml round bottom flask and 25 ml, 0.5 N alcoholic potassium hydroxide solution added in same flask. The flask was fitted with a long air condenser and heated solution at reflux temperature about 30 minutes. Finally, the flask was cooled and added 1 ml of 1 percent phenolphthalein solution and titrate the excess of the alkali against standard 0.5 N HCl. At the same time and under similar conditions carry out a blank titration without fat (25 ml, 0.5 N same alcoholic KOH solution was taken in another round bottom flask and heated in a similar way and titrated, against 0.5 N acid) [14-17].

Iodine value

5 g of oil was taken into 200 ml a glass stoppard bottle. 5 ml of CCl₄ was added to dissolve this oil after 25 ml of Wij's solution was added and to allow it at least 1 hour in a dark place. Then 5 ml of 10% potassium iodide solution and 50 ml water were added to each bottle and titrated against 0.1N Na₂S₂O₃ solution using starch solution as the indicator, near the end point of titration the color of solution becomes pale yellow. Blue color disappears which indicates the end point. At the same time and under similar conditions carry out a blank titration without oil [14-17].


Peroxide Value

The peroxid value is a measure of the concentration of substances that oxidize potassium iodide to iodine. 5 g weight of oil samples is dissolved in acetic acid then chloroform and saturated KI mixture are added to the sample and the amount of iodine liberated from KI by the oxidative action of peroxides present in the oil is determined by titration with 0.1 N sodium thiosulphate using starch solution as an indicator. Titration was also performed for blanks [14-17].

Preparation of fatty acid methyl esters (FAMES):

For conversion of corn oil into FAME the following methods was adopted: around 0.2 gm corn oil was taken into 50 mL glass vial

then mix with 5 mL of 0.50 N methanolic NaOH, the mixture was heated for 4 mins at 60 C. The mixture was allowed to cool at room temperature, then 6 mL of 14% BF₃ solution was added [11] to the mixture and again heated for 3 mins at 60 C. The mixture was again cool at room temperature then added 10 mL , isooctane and shake it well, then keep it to settling down. After settling the mixture, the upper layer of the mixture was transferred to the tube containing sodium sulfate to remove the moisture. The extract was analyzed with GC-FID (Nuckon - 5700), the flow chart of the corn oil analysis method condition shown in Scheme 1 [12].



Scheme 1. Flow chart of the corn oil fatty acid analysis.

Chromatography

Nuckon – 5700 gas chromatograph equipped with a flame-ionization detector (FID) and glass column, digital controls, separate independent proportional temperature controllers for oven, detector, Injector, Detector temperature 390 0 C in 10 C steps set by thumbwheel switches, oven temperature 399 0 C in 1 C step set by thumbwheel switches. Temperature programming: - microprocessor based programmer with rate 0.1 0 C – 29.9 0 C per minute with initial time hold up to 99 minute. Dual channel data station, 2 detectors can be operated simultaneously. Use Nitrogen gas cylinder -60 psi. Hydrogen gas cylinder -20 psi. Zero air gas cylinder -20 psi.

Certified Reference Materials (CRM)

Sigma, Catalog no 46961, Lot no - LC 26752V, Exp. May 2020.

Results

Corn oil show physical properties liquid foam, yellow colour, characteristic odour, slightly soluble in alcohol and miscible in ether, chloroform, petroleum ether, benzene, hexane and other chemical solvents (table -1).

Table 1: Physical properties of corn oil

Property	Characteristic/ Value
Physical state at 25 ⁰ C/ 1Atm	Liquid
Colour	Yellow
Odor	Characteristic Corn
Solubility	Slightly soluble in alcohol
Miscibility	Miscible in ether, chloroform, petroleum ether, benzene, hexane

The quality of Corn oil was analyzed by evaluating physicochemical properties such as density /specific gravity, viscosity, flash point, boiling point, saponification value, unsaponifiable value, iodine value, colour (Lovibond tintometer),

peroxide value and refractive Index. Results are presented in Table 2.

Table 2: Physicochemical properties of corn oil

Property	Characteristic/ Value
Density (g/mL)/specific gravity	0.918 at 25°C
Viscosity (millipoise)	114.00
Flash point	305
Boiling Point (°C)	148.00
Saponification Value (mg)	153.6
Unsaponifiable value in gm	2.2
Iodine Value (g)	15.77
Colour (Lovibond tintometer)	3.0 red max.
Peroxide value (meq/kg)	0.165
Refractive Index (40° C)	1.466

Corn seed samples collected from the local market alwar, rajasthan were analyzed on the composition of fatty acids using gas chromatographic FAMES method. The content of following saturated and unsaturated fatty acids was tested in the samples: palmitic acid, stearic acid, oleic acid, lenoleic acid, linolenic acid and arachidic acid. The fatty acid percent compositions of corn oil are shown in table 3 to 4 and figure 1 respectively.




Fig 1: Gas chromatogram of corn oil

Table 3: Presenting the different parameters of the fatty acid of the corn oil.

Peak No.	RT (Min)	Area (mV-Sec)	Height (mVolt)	RF	Amount (ML)	Amount%	Component Name
1	13:34	317.555	19.153	1.000	317.555	10.386	Palmitic
2	16:40	145.137	6.903	1.000	145.137	4.747	Stearic
3	17:16	1,256.088	54.738	1.000	1,256.088	41.081	Oleic
4	18:15	1,309.805	69.792	1.000	1,309.805	42.838	Lenoleic
5	19:6	24.402	1.668	1.000	24.402	0.798	Linolenic
6	19:39	4.568	0.365	1.000	4.568	0.149	Arachidic

Table 4: The content of SFA, MUFA, PUFA (% w/w)

Saturate d fatty acid	Monounsaturate d fatty acid	Polyunsaturate d fatty acid	unsaturate d fatty acid
15.282	41.081	43.636	84.717

Unsaturated fatty acids are classified as either monounsaturated fatty acids, because they have only one double bond (e.g., omega-7 and -9 fats), and polyunsaturated fatty acids, since they have more than one double bond in their backbone (e.g., omega-3 and -6 acids) [25, 26]. Saturated fatty acids are in solid state at room

temperature. The main sources include ghee, butter, tallow coconut, lard, palm, and palm kernel oil. Saturated fatty acids are also called nonessential fats (not to be confused as unimportant) [27-31].

Conclusion

The observation results concluded that we should consider the physical, physicochemical properties and fatty acid profile for selecting good type corn oil.

Competing Interests

Authors have declared that no competing interests exist.

Acknowledgement

The author is thankful to the Nirmal Industries Pvt Ltd, Alwar for providing the all facilities as well as the instrument and chemical for the Content of Fatty Acid in Corn Seed Oil using GC-FID method.

References

1. Staller JE, Tykot R, Benz B. Histories of Maize: Multidisciplinary Approaches to the Prehistory, Linguistics, Biogeography, Domestication, and Evolution of Maize. Walnut Creek: Left Coast Press, 2006.
2. Sofi PA, Wani SA, Rather AG, Wani SH. Review article: Quality protein maize (QPM): Genetic manipulation for the nutritional fortification of maize. *J Plant Breed Crop Sci.* 2009; 1(6):244-53.
3. Vasal SK. The quality protein maize story. *Food Nutr Bull.* 2000; 21(4):445-50.
4. Alexander DE. Breeding special nutritional and industrial types. In: *Corn and Corn Improvement* (Eds. Sprague, G. F. and Dudley, J. W.). ASA, CSSA and SSSA, Madison, Wisconsin, 1988, 869-80.
5. Alexander DE, Lambert RJ. Relationship of kernel oil content to yield in maize. *Crop Sci.* 1968; 8:273-74.
6. Alrefai R, Berke TG, Rocheford TR. Quantitative trait locus analysis of fatty acid concentrations in maize. *Genome.* 1995; 38:894-901.
7. Batista LAR, Tosello GA. Influência da fonte polinizadora sobre o conteúdo de óleo em grãos de milho. *Pesquisa Agropecuária Brasileira.* 1982; 17:1757-62.
8. Bergquist RR, Nubel DS, Thompson DL. 1998. Production method for high oil corn grain. US Patent Number: 5704160.
9. Gutiérrez LF, Rosada LM, Jiménez A. Chemical composition of Sacha inchi (*Plukenetia volubilis* L.) seeds and characteristics of their lipid fraction. *Grasas Aceites* 2011; 62(1):76-83.
10. Popa VM, Gruia A, Raba DN, Dumbrava D, Moldovan C, Bordean D, Mateescu C. Fatty acids composition and oil characteristics of linseed (*Linum usitatissimum* L.) from Romania. *J. Agroalim. Processes Technol.* 2012; 18(2): 136e140.
11. Piskernik S, Vidrih R, Demsar L, Koron D, Rogelj M, Zontar TP. Fatty acid profiles of seeds from different Ribes species. *LWT - Food Sci. Technol.* 2018; 98:424e427.
12. Manual of method of analysis of foods. Food safety and standards authority of India, ministry of health and family welfare government of India, new delhi. pp.3-88, 2015

13. Sharif RH, Paul RK, Bhattacharjya DK, Ahmed KU. "Physicochemical characters of oilseeds from selected mustard genotypes" Physicochemical characters of oilseeds. 2017; 15(1):27-40.
14. Purendra Singh. Physico-chemical investigations of Mustard seed (*Brassica juncea* L), International Journal of Scientific Research in Multidisciplinary Studies. 2018; 4(6):24-27.
15. Hasan Ms, Jahan R, Alam MA, Khatun MK, SMA Reza, "Study on Physicochemical Properties of Edible Oils Available in Bangladeshi Local Market" Archives of Current Research International. 2016; 6(1):1-6.
16. Purendra Singh, Hand book of phytochemical, pharmacognostical, water and edible oil analysis (A lab manual), Educreation Publishing, 2019, 1-96.
17. Sabate J. Nut composition and body weight. American Journal of Clinical Nutrition. 2003; 78:647s-650s
18. Willett WC. Is dietary fat a major determinant of body fat? American Journal of Clinical Nutrition. 1998; 67:556s-62s.
19. Anthea M, Hopkins J, McLaughlin CW, et al. Human Biology and Health, Prentice Hall, Englewood Cliffs, NJ, USA, 1993.
20. Mendil D, Uluözlü OD, Tuzen M, Soylak M. Investigation of the levels of some element in edible oil samples produced in Turkey by atomic absorption spectrometry," Journal of Hazardous Materials. 2009; 165:724-728,. View at Google Scholar
21. Johannesen E. Palm fruit oil-much more than ordinary oil," Journal of Science in Africa Magazine, 2005.
22. Manorama R, Rukmini C. Nutritional evaluation of crude palm oil, The Oil Technologists' Association of India. 1991; 22:83-87,. View at Google Scholar
23. Shaik R. Analytical tool for rapid analysis of edible oil [M.S. thesis], Aalborg University Esbjerg, 2008.
24. Patkowska-Sokoła B., Usydus Z., Szlinder-Richert J., Bodkowski R. Technology for recovering omega-3 fatty acids from family from fish oils and protecting them against oxidative changes. Przem. Chem. 2009; 88:548-552. [Google Scholar]
25. Meyer B.J., Neil J., Mann N.J., Lewis J.L., Milligan G.C., Sinclair A.J., Howe P.R.C. Dietary intakes and food sources of omega-6 and omega-3 polyunsaturated fatty acids. Lipids. 2003; 38:391-398. doi: 10.1007/s11745-003-1074-0. [PubMed] [CrossRef] [Google Scholar]
26. Willett WC. Dietary fats and coronary heart disease. J Intern Med 2012; 272:13-24.
27. Bier DM. Saturated fats and cardiovascular disease: Interpretations not as simple as they once were. Crit Rev Food Sci Nutr 2016; 56:1943-6.
28. Praagman J, Beulens JW, Alsema M, Zock PL, Wanders AJ, Sluijs I, et al. The association between dietary saturated fatty acids and ischemic heart disease depends on the type and source of fatty acid in the European Prospective Investigation into Cancer and Nutrition-Netherlands cohort. Am J Clin Nutr 2016; 103:356-65.
29. Mead, JF, Alfin-Slater RB, Howton DR, Popják G. Lipids: Chemistry, Biochemistry and Nutrition. New York: Plenum Press; 1986.
30. Blackburn GL, Kater G, Mascioli EA, Kowalchuk M, Babayan VK, kBistrián BR. A reevaluation of coconut oil's effect on serum cholesterol and atherogenesis. The Journal of the Philippine Medical Association 1989; 65:144-152.